

## Original Article

**Cite this article:** Rhead R, MacManus D, Jones M, Greenberg N, Fear NT, Goodwin L (2022). Mental health disorders and alcohol misuse among UK military veterans and the general population: a comparison study. *Psychological Medicine* 52, 292–302. <https://doi.org/10.1017/S0033291720001944>

Received: 10 June 2019

Revised: 19 May 2020

Accepted: 21 May 2020

First published online: 10 August 2020

**Key words:**


Alcohol misuse; anxiety; common mental disorder; depression; military; veteran

**Author for correspondence:**

Rebecca Rhead,

E-mail: [rebecca.rhead@kcl.ac.uk](mailto:rebecca.rhead@kcl.ac.uk)

# Mental health disorders and alcohol misuse among UK military veterans and the general population: a comparison study

Rebecca Rhead<sup>1,2</sup> , Deirdre MacManus<sup>1,2</sup>, Margaret Jones<sup>2</sup>, Neil Greenberg<sup>2</sup>, Nicola T Fear<sup>2,3</sup> and Laura Goodwin<sup>2,4</sup>

<sup>1</sup>Department of Forensic and Neurodevelopmental Science, King's College London, London, UK; <sup>2</sup>King's Centre for Military Health Research, Department of Psychological Medicine, King's College London, London, UK; <sup>3</sup>Academic Department of Military Mental Health, Department of Psychological Medicine, King's College London, London, UK and <sup>4</sup>Department of Psychological Sciences, University of Liverpool, Liverpool, UK

**Abstract**

**Background.** For a small minority of personnel, military service can have a negative impact on their mental health. Yet no studies have assessed how the mental health of UK veterans (who served during the recent operations in Afghanistan or Iraq) compares to non-veterans, to determine if they are at a disadvantage. We examine the prevalence of mental disorders and alcohol misuse in UK veterans compared to non-veterans.

**Methods.** Veteran data were taken from the third phase of the King's Centre for Military Health Research cohort study ( $n = 2917$ ). These data were compared with data on non-veterans taken from two large general population surveys: 2014 Adult Psychiatric Morbidity Survey ( $n = 5871$ ) and wave 6 of the UK Household Longitudinal Study (UKHLS,  $n = 22\,760$ ).

**Results.** We found that, overall, UK veterans who served at the time of recent military operations were more likely to report a significantly higher prevalence of common mental disorders (CMD) (23% *v.* 16%), post-traumatic stress disorder (PTSD) (8% *v.* 5%) and alcohol misuse (11% *v.* 6%) than non-veterans. Stratifying by gender showed that the negative impact of being a veteran on mental health and alcohol misuse was restricted to male veterans. Being ill or disabled was associated with a higher prevalence of CMD and PTSD for both veterans and non-veterans.

**Conclusion.** Whilst the same sociodemographic groups within the veteran and non-veteran populations seemed to have an increased risk of mental health problems (e.g. those who were unemployed), male veterans, in particular, appear to be at a distinct disadvantage compared to those who have never served.

**Introduction**

The majority of UK Armed Forces (AF) personnel cope well with the challenges of military service and successfully adapt to veteran life after they leave (Royal British Legion, 2014). For some, military service can negatively impact their mental health (Iversen & Greenberg, 2009; Stevelink et al., 2018). Ensuring that the mental health needs of veterans are met is a UK Government priority (Defence Select Committee, 2018) and legal obligation is enshrined in the AF Covenant (Taylor, 2011). Despite this, there is little UK research into how the mental health of veterans compares to the general population.

Much international research has been conducted on the mental health of *servicing* military personnel (Fear et al., 2010; Hooff et al., 2014; Lee, Sudom, & Zamorski, 2013; Sundin et al., 2014). UK comparisons made between *servicing* personnel and working non-veterans found symptoms of common mental disorders (CMD) to be more common among *servicing* military personnel (Goodwin et al., 2015). Alcohol misuse has also been found to be more common in UK military personnel than the general population with heavy drinkers less likely to change their drinking patterns over time (Fear et al., 2007; Goodwin et al., 2017; Thandi et al., 2015).

Though studies from other countries have shown veterans to have poorer mental health than non-veterans (Hoerster et al., 2012; Oster, Morello, Venning, Redpath, & Lawn, 2017), findings from the few UK studies comparing military veterans and non-veteran mental health have suggested that there is little difference between the two groups (Murphy, Palmer, Westwood, Busuttill, & Greenberg, 2016; Woodhead et al., 2011a, b). Analysis restricted to a representative sample of post-national service veterans ( $n = 257$ ) found no statistically significant difference in the prevalence of poor mental health between veterans and non-veterans (Woodhead et al., 2011b). However, the study utilised a small sample of veterans, few of whom had served while recent military operations were ongoing in Iraq and Afghanistan. Recent estimates suggest that probable post-traumatic stress disorder (PTSD) is now higher

among UK veterans compared to those in service (Stevellink et al., 2018). Whether it is also higher compared to the general population remains uncertain, as is our understanding of the manifestation of PTSD; veterans may experience more severe or differing symptoms than non-veterans (Vasterling et al., 2010; Wilkins, Lang, & Norman, 2011).

This study compares the prevalence of CMD, PTSD and alcohol misuse, and the associated domains of PTSD and alcohol misuse, in current cohorts of veterans who served at the time of recent conflicts and in non-veterans using large, nationally representative samples. Socio-demographic characteristics such as employment status and socio-economic grade are commonly associated with mental health (Ostler et al., 2001), these factors will be adjusted for in our analysis and prevalence estimates will be made in working and non-working groups as well as overall. Analyses will also examine if there are gender differences in how veterans compare to equivalent non-veterans.

This study aims to:

- (1) Establish how the prevalence of mental health problems and domains compare for veterans and non-veterans, by gender, including individual comparisons for working and non-working groups.
- (2) Examine the association of socio-economic grade, employment status, education and marital status with mental health, and how these associations differ between veterans and non-veterans.

## Methods

### Data

#### Veterans

Our veteran sample was taken from phase 3 of the King's Centre for Military Health Research (KCMHR) cohort study, a large ongoing study of UK military personnel assessing the physical and mental health consequences (CMD, PTSD and alcohol misuse) of military service and deployment. Previous waves of the cohort were collected between June 2004 and March 2006 (phase 1) (Hotopf et al., 2006), and November 2007 and September 2009 (phase 2) (Fear et al., 2010). This study uses the most recent phase (phase 3) for which data was collected between October 2014 and December 2016 with an overall response rate of 44% (Stevellink et al., 2018). Sample weights were calculated as the inverse probability of an individual being sampled from a specific subpopulation (TELIC, Era, HERRICK, phase two replenishment, phase three replenishment) and specific engagement type (regular or reserve). Response weights were generated to account for non-response and calculated as the inverse probability of responding once sampled, driven by factors shown empirically to predict the response (gender, age, rank, service, engagement type, serving status, subpopulation and the interaction between subpopulation and engagement type). The sample weights and response weights were multiplied together to make a single weight. The weighted analyses provided valid results under the assumption that the data were missing at random and that the observed variables modelled to drive non-response were correctly identified (see (Stevellink et al., 2018)).

This study defines veterans as those who have served in the military as a regular and have since left. Those still engaging in active service, and those who *only* served as reservists were excluded from these analyses.

#### Non-veterans (general population)

Our non-veteran sample was taken from two surveys; the 2014–2016 UK Household Longitudinal Study (UKHLS) and the 2014 Adult Psychiatric Morbidity Survey (APMS). The use of two surveys was necessary as mental health assessments needed to be made using the same measures of PTSD, CMD and alcohol misuse as used by the KCMHR cohort study to ensure comparability – UKHLS contains the same measure of CMD, while APMS contains the same measures of PTSD and alcohol misuse. No single general population survey with all three measures was available.

The UKHLS is the largest panel survey in the world. Launched in 2008 as a continuation of the British Household Panel Survey (Lynn, 2006), the UKHLS follows the original sample and also includes a newly-selected General Population Sample (GPS) of 25 500 responding households across the UK. The GPS is based upon a proportionately stratified, equal probability (clustered) sample of residential addresses drawn to a uniform design throughout England, Scotland and Wales (Lynn, 2006). Wave 6 of UKHLS (2014–2016) achieved an individual adult response rate of 65%.

The APMS provides data on the prevalence of both treated and untreated psychiatric disorders in the English adult population (aged 16 yrs and over). The sample for APMS 2014 was designed to be representative of the population living in private households in England, adopting a multi-stage stratified probability sampling design (McManus, Bebbington, Jenkins, & Brugha, 2016) and achieved a response rate of 57%. Though both datasets included optional weights, neither were used in this analysis.

### Measures

#### Mental health and alcohol misuse

Mental health and alcohol misuse were evaluated using the 12-item GHQ-12 (Goldberg et al., 1997) to assess CMD, the 17-item National Centre for PTSD Checklist – Non-veteran version (PCL-C) (Weathers, Litz, Herman, Huska, & Keane, 1994) to assess PTSD, and the 10-item World Health Organization's Alcohol Use Disorders Identification Test (AUDIT) to identify probable alcohol misuse (Babor, Higgins-Biddle, Saunders, & Monteiro, 2001; Fear et al., 2007).

GHQ-12 symptoms were rated on a four-point scale. A bi-modal scoring method of 0–0–1–1 was used (Goldberg et al., 1997). Possible scores ranged from 0 to 12 with a cut-off of 4 or more to represent probable CMD (Goldberg et al., 1997). PCL-C symptoms were rated 1–5. Possible scores ranged from 17 to 85 with a cut-off point of 50 or more to assess caseness (Weathers et al., 1994). DSM-IV domains: re-experiencing, avoidance and hyperarousal were also assessed (Taylor, Koch, Kuch, Crockett, & Passey, 1998). The study is restricted to the use of DSM-IV criteria as PCL-C was the only measure of PTSD included in APMS.

AUDIT scores range from 0 to 40 and a cut off of 16 or more was used to assess caseness (Babor et al., 2001; Fear et al., 2007). The AUDIT's three mutually exclusive domains were also examined (Babor et al., 2001):

- **Hazardous drinking:** a pattern of alcohol consumption that increases the risk of harmful consequences for the user or others – *based on frequency and quantity of alcohol consumption.*
- **Evidence of present harm:** alcohol consumption that results in consequences to physical and mental health – *indicated by drinking-related injuries, blackouts and/or guilt as well as concern from others.*

- **Dependant drinking:** a strong desire to consume alcohol and diminished control over its use despite harm-full consequences – indicated by *impaired control over drinking, increased salience of drinking and drinking in the morning.*

### Socio-demographic characteristics

The socio-demographic characteristics of veterans and non-veterans were examined using data on gender, age in years (22–34, 35–44, 45–54, 55–64, 65–74), level of education (no qualification, GCSE or equivalent, A-level or degree level), marital status (married/in a relationship, single or divorced/separated/widowed), and employment status (employed part-time/full-time, unemployed, retired, long-term ill/disabled or economically inactive).

A dichotomised version of employment status was used in our comparison of mental health prevalence (to assess mental health differences by veteran status in working and non-working groups) with employed part-time/full-time defined as working, and all other responses defined as non-working.

A five-category NS-SEC social grade variable (Office for National Statistics, n.d.) was available in both the APMS and UKHLS datasets. This data was not available in the KCMHR cohort and as such, veterans were manually assigned an NS-SEC social grade based on their reported job title. A total of 6.1% of employed veterans were unable to assign an NS-SEC social grade as their job title was not reported ( $n = 195$ ), or their job title was not sufficiently detailed to allow for a classification ( $n = 223$ ).

### Statistical analysis

#### Restricting the military sample

Of the 8094 personnel who participated in phase 3 of the KCMHR cohort study, 3008 met our definition of a veteran (a former regular who is no longer affiliated with the military). Thirty-five veterans were excluded as information on employment status was not provided. Further exclusions were made during mental health comparative analysis as some veterans did not respond to any items on the respective scale (GHQ  $n = 56$ , PCL-C  $n = 67$ , AUDIT  $n = 64$ ). Our veteran sample size ranged from 2906 to 2917 depending on the mental health outcome. The sample size was reduced ( $n = 418$ ) for models where social grade (NS-SEC) of employed veterans was accounted for.

#### Matching samples on age and gender

To enable comparisons between veterans and non-veterans, our non-veteran samples were reweighted to match on age and gender with the veteran sample.

For our comparison of CMD across veterans and non-veterans, KCMHR and UKHLS data were combined into a single dataset. UKHLS participants outside the age range of our veteran sample (22–74 years) were removed from the sample ( $n = 8065$ ) as were those who reported currently or previously serving in the AF ( $n = 75$ ). The remaining sample of 22 760 was reweighted to match the exact age and gender of veterans in the KCMHR data using entropy balancing [a multivariate reweighting method described in (Hainmueller, 2012)] to ensure comparability.

Entropy balancing allows the researcher to obtain a high degree of covariate balance between samples, it is also versatile and retains valuable information in the pre-processed data by allowing the unit weights to vary smoothly across units. Entropy balancing is only problematic if misused. Imposing a large number of constraints or setting unrealistic balance constraints with small and/or limited data can lead to inflated standard errors. To avoid this, we have

only imposed two constraints (age and gender) and have used large datasets without unrealistic balance constraints.

Separate analyses were conducted for comparisons of PTSD and alcohol misuse in veterans and non-veterans where KCMHR and APMS data were combined into a single dataset. APMS participants outside the age range of the veteran sample ( $n = 1157$ ) or who reported ever serving in the AF ( $n = 518$ ) were excluded. The remaining sample of 5871 was re-weighted to match on age and gender using entropy balancing.

For each set of analysis (KCMHR v. UKHLS & KCMHR v. APMS) our veteran sample was weighted for non-response (Stevellink et al., 2018), and entropy weights were applied to the respective non-veteran samples to ensure age and gender comparability. All proportions and odds ratios reported in this study are weighted, frequencies are unweighted.

### Analysis

- (1) Prevalence of GHQ-12, PCL-C and AUDIT caseness were calculated, and adjusted logistic regression analyses used to compare veterans and non-veterans in working and non-working groups after accounting for socio-demographic characteristics (age, gender, education, marital status, employment and social grade). These comparisons were stratified by gender, as gender moderated the effect of veteran status on mental health caseness.
- (2) Caseness for AUDIT and PCL-C domains were compared between veterans and non-veterans using unadjusted and logistic regressions (adjusting for socio-demographic characteristics).
- (3) Using logistic regression, the association between socio-demographic characteristics and mental health caseness was assessed in veteran and non-veteran samples separately. Interaction effects were tested using a Wald test to compare models with and without an interaction term to determine whether sociodemographic characteristics moderated the effect of being a veteran (compared to not being a veteran) on mental health outcomes.
- (4) Sensitivity analyses were conducted in males and females separately to examine differences between veterans and non-veterans, stratifying the samples based upon the veterans' military experience. The veteran sample was stratified by i) deployment to Iraq or Afghanistan (yes, no) and ii) restricted to only those who were deployed, whether they had deployed in a combat role (yes, no) (in males only). Veterans within each of these strata were matched on age and gender to non-veterans, and the prevalence of probable CMD, PTSD and alcohol misuse was compared using logistic regression before and after adjusting for education and marital status (as indicators of socio-economic status and a proxy for social support).

All data were analyzed using the statistical software package Stata (version 14).

### Results

Table 1 shows the sociodemographic characteristics for KCMHR, APMS and UKHLS samples after entropy balancing.

#### Mental health prevalence and employment status

Overall, veterans report significantly higher prevalence of CMD (23% v. 16%), PTSD (8% v. 5%) and alcohol misuse (11% v. 6%) than non-veterans (see Table 2). Caseness for CMD and

**Table 1.** Sociodemographic characteristics for veteran and general population samples

	KCMHR (N = 2917)		APMS (N = 5871)		UKHLS (N = 22 760)	
	N	%	N	%	N	%
<i>Sociodemographic characteristics</i>						
Gender						
Male	2612	91.4%	2259	90.9%	9950	91.4%
Female	305	8.6%	3612	9.1%	12 810	8.6%
Age (yrs)						
22-34	453	35.9%	1215	41.7%	5238	39.7%
35-44	820	27.3%	1147	26.3%	4742	23.2%
45-54	1091	26.7%	1232	15.7%	5243	19.2%
55-64	472	8.8%	1155	10.2%	4289	11.3%
65-74	81	1.4%	1122	6.1%	3248	6.5%
Marital status						
Married/long-term relationship	2512	84.8%	3554	60.7%	16 055	68.1%
Single	190	9.3%	1181	30.2%	3994	26.2%
Separated, divorced or widowed	198	5.9%	1136	9.2%	2641	5.7%
Highest qualification						
No qualifications	130	4.5%	1147	15.0%	1976	5.3%
GCSE of equivalent	659	24.3%	1597	26.0%	5581	23.1%
A level	991	39.3%	972	20.6%	4653	24.9%
Degree or higher	1133	32.0%	2097	38.5%	10 550	46.7%
Employment status						
In employment (ft/pt)	2567	90.7%	3588	76.2%	15 367	78.5%
Unemployed (looking for work)	85	3.0%	177	4.1%	1036	6.4%
Long-term ill or disabled	67	2.2%	341	6.4%	760	2.8%
Retired	172	3.1%	1192	6.8%	3707	7.5%
Economically inactive	26	1.0%	573	6.5%	1890	4.8%
NSSEC grade (employed only)						
1: Managerial and Professional	843	32.9%	1614	47.3%	6891	46.2%
2: Intermediate	522	25.1%	578	9.8%	2014	10.1%
3: Small and own employers	105	5.4%	376	11.2%	1488	11.0%
4: Lower supervisory & technical	165	9.4%	164	7.6%	1031	9.4%
5: Semi-routine/routine	516	27.1%	842	24.1%	3598	23.3%

Sample size may vary slightly due to missing data in outcome variables.

PTSD is also significantly lower for working people compared to non-working people for both veterans and non-veterans.

Amongst men, being a veteran was significantly associated with reporting probable CMD for both working and non-working groups. The strength and significance of these associations remain largely unchanged when adjusting for socio-demographic characteristics. There was no significant difference in CMD prevalence between female veterans and female non-veterans regardless of the working group.

Male veterans in both working and non-working groups were significantly more likely to report probable PTSD than non-veterans. Female veterans in employment were significantly more likely to report probable PTSD than employed women

from the general population (there was no significant difference amongst non-working women).

Only employed male veterans were significantly more likely to report alcohol misuse than their non-veteran counterparts. Being a veteran was not found to be associated with reporting alcohol misuse for women, regardless of the working group.

### Domains

Male veterans were significantly more likely to meet criteria for all PTSD domains. There was no statistically significant association between being a veteran and any PTSD domain for women. Male veterans were significantly more likely to meet the criteria

**Table 2.** Sample on mental health caseness by employment status

		Veteran		Non-veteran		Crude		Adjusted for marital status and education		Adjusted for marital status, education, employment status and NS-SEC grade	
		n	%	n	%	OR	CI	OR	CI	OR	CI
Overall											
CMD caseness	Full sample	618	23.3%	4063	16.0%	1.59	1.40,1.81	1.67	1.46,1.91	1.84	1.60,2.12
	Working	507	21.5%	2329	13.3%	1.79	1.55,2.06	1.89	1.63,2.20	1.86	1.58,2.18
	Non-working	111	40.4%	1734	25.9%	1.94	1.44,2.62	1.65	1.18,2.31	1.59	1.09,2.30
PTSD caseness	Full sample	187	7.9%	293	5.2%	1.57	1.19,2.07	2.29	1.67,3.14	2.77	1.95,3.94
	Working	143	6.7%	100	3.2%	2.16	1.47,3.19	2.74	1.81,4.13	2.76	1.82,4.20
	Non-working	44	19.2%	193	11.8%	1.78	1.11,2.84	2.45	1.34,4.47	2.86	1.49,5.50
Alcohol misuses (16+)	Full sample	264	10.7%	175	5.7%	1.98	1.51,2.59	2.30	1.73,3.07	2.32	1.74,3.10
	Working	228	10.6%	103	5.1%	2.20	1.60,3.03	2.48	1.79,3.44	2.59	1.84,3.65
	Non-working	36	11.6%	72	7.8%	1.56	0.90,2.70	1.66	0.91,3.01	1.40	0.75,2.61
Men											
CMD caseness	Full sample	557	23.3%	1466	15.5%	1.65	1.45,1.89	1.73	1.50,2.00	1.93	1.66,2.25
	Working	459	21.7%	905	13.0%	1.86	1.60,2.16	1.98	1.68,2.32	1.96	1.65,2.33
	Non-working	98	39.7%	561	25.6%	1.95	1.41,2.70	1.55	1.07,2.24	1.64	1.09,2.47
PTSD caseness	Full sample	169	7.9%	101	5.0%	1.63	1.20,2.21	2.34	1.66,3.29	2.86	1.95,4.21
	Working	130	6.8%	40	3.2%	2.21	1.46,3.37	2.79	1.79,4.35	2.84	1.80,4.46
	Non-working	39	19.0%	61	11.5%	1.81	1.08,3.03	2.40	1.22,4.70	3.09	1.45,6.58
Alcohol misuses (16+)	Full sample	253	11.4%	106	6.1%	1.99	1.51,2.63	2.33	1.73,3.13	2.36	1.75,3.19
	Working	220	11.3%	65	5.4%	2.24	1.61,3.12	2.52	1.80,3.53	2.63	1.85,3.74
	Non-working	33	12.4%	41	8.5%	1.51	0.85,2.69	1.60	0.85,3.00	1.37	0.71,2.66
Women											
CMD caseness	Full sample	61	23.1%	2597	21.2%	1.12	0.80,1.56	1.20	0.86,1.69	1.12	0.80,1.56
	Working	48	19.6%	1424	18.2%	1.15	0.79,1.66	1.21	0.84,1.74	1.14	0.78,1.66
	Non-working	13	45.4%	1173	27.9%	1.88	0.85,4.18	1.91	0.91,4.02	0.75	0.38,1.48
PTSD caseness	Full sample	18	7.6%	192	6.8%	1.13	0.63,2.03	2.07	1.02,4.20	1.75	0.93,3.30
	Working	x	x	60	3.5%	1.65	0.83,3.31	2.21	1.00,4.89	2.40	1.04,5.57
	Non-working	x	x	132	13.8%	1.62	0.52,5.02	2.15	0.77,6.03	0.72	0.24,2.18
Alcohol misuses (16+)	Full sample	11	3.3%	69	2.2%	1.49	0.74,3.01	1.56	0.80,3.06	1.57	0.76,3.23
	Working	x	x	38	2.0%	1.41	0.61,3.26	1.47	0.63,3.47	1.80	0.79,4.12
	Non-working	x	x	31	2.6%	2.46	0.68,8.94	2.25	0.63,8.05	2.46	0.54,11.13

x = Suppressed due to low *n* (threshold is 10).

a = NSSEC grade only adjusted for in working groups.

for hazardous, dependent and harmful drinking than non-veterans. Female veterans were significantly more likely to report symptoms of hazardous drinking only (compared to female non-veterans) (see Table 3).

#### *Sociodemographic characteristics associated with poor mental health*

Table 4 shows the age and gender-adjusted associations between socio-demographic characteristics and mental health caseness for veterans and non-veterans separately.

#### *Common mental disorders*

Employment status ( $p < 0.01$ ,  $\chi^2 = 12.9$ ,  $df = 4$ ), gender ( $p < 0.005$ ,  $\chi^2 = 9.2$ ,  $df = 1$ ) and age ( $p < 0.05$ ,  $\chi^2 = 11.1$ ,  $df = 4$ ) were found to significantly moderate the effect of being a veteran on CMD.

Being unemployed and long-term ill or disabled was associated with CMD caseness for both veterans and non-veterans, although the size of the effect was larger for veterans. Amongst the general population, gender was significantly associated with probable CMD, with women having greater odds of meeting caseness criteria. There was no statistically significant association between gender and CMD caseness amongst veterans. Veterans aged 45+

**Table 3.** Mental health domain prevalence in veterans and non-veterans

		Veteran		Non-veteran		Crude		Adjusted for marital status and education		Adjusted for age, marital status, education, employment status and NS-SEC grade	
		N	%	N	%	OR	CI	OR	CI	OR	CI
<b>Men</b>											
PCL-C domains	<i>Re-experiencing</i>	492	11.7%	417	9.5%	1.24	1.04,1.49	1.58	1.29,1.93	1.77	1.39,2.25
	<i>Avoidance</i>	406	9.1%	231	5.6%	1.74	1.42,2.15	2.32	1.82,2.96	2.69	2.01,3.60
	<i>Hyperarousal</i>	544	12.5%	373	9.3%	1.44	1.21,1.73	1.86	1.52,2.28	2.20	1.74,2.79
AUDIT domains	<i>Hazardous</i>	2323	51.2%	1340	36.1%	4.03	3.23,5.03	4.24	3.34,5.37	5.00	3.70,6.75
	<i>Present Harm</i>	1107	30.1%	686	21.5%	1.44	1.21,1.73	1.86	1.52,2.28	2.20	1.74,2.79
	<i>Alcohol Dependence</i>	614	17.9%	320	11.1%	1.74	1.42,2.15	2.32	1.82,2.96	2.69	2.01,3.60
<b>Women</b>											
PCL-C domains	<i>Re-experiencing</i>	53	10.6%	863	13.5%	0.78	0.54,1.12	0.96	0.66,1.41	0.85	0.55,1.31
	<i>Avoidance</i>	38	7.9%	403	6.6%	1.31	0.86,2.00	1.93	1.17,3.19	1.38	0.80,2.39
	<i>Hyperarousal</i>	52	9.6%	693	10.8%	0.91	0.63,1.31	1.09	0.74,1.60	1.02	0.66,1.57
AUDIT domains	<i>Hazardous</i>	228	42.4%	1560	30.3%	1.69	1.19,2.40	1.66	1.13,2.44	1.78	1.17,2.72
	<i>Present Harm</i>	84	19.7%	762	16.5%	0.91	0.63,1.31	1.09	0.74,1.60	1.02	0.66,1.57
	<i>Alcohol Dependence</i>	43	9.2%	309	7.0%	1.31	0.86,2.00	1.93	1.17,3.19	1.38	0.80,2.39

years were less likely to meet the threshold for CMD caseness than younger participants, though, for non-veterans, only those aged 65+ years were significantly less likely to indicate probable CMD.

### PTSD

Marital status ( $p < 0.01$ ,  $\chi^2 = 8.20$ ,  $df = 2$ ) and age ( $p < 0.01$ ,  $\chi^2 = 15.42$ ,  $df = 4$ ) were found to significantly moderate the effect of being a veteran on PTSD. For both samples, being single was significantly associated with probable PTSD compared to being in a relationship, although the association was significantly smaller for veterans. Veterans aged 44–74 yrs were less likely to meet criteria for PTSD caseness than participants aged 22–44 yrs, as were non-veterans aged 65–74 yrs.

Although no statistically significant interactions were found for level of education or employment status, having any educational qualification was associated with a decreased likelihood of meeting criteria for PTSD caseness for both samples compared to having no qualifications. Additionally, for both samples, compared to being employed, being unemployed and long-term ill or disabled was associated with PTSD caseness.

### Alcohol misuse

Marital status significantly moderated the effect of being a veteran on alcohol misuse ( $p < 0.01$ ,  $\chi^2 = 14.26$ ,  $df = 2$ ). For non-veterans, being single (compared to being in a relationship) was associated with alcohol misuse. For veterans, being separated, divorced or widowed (compared to being in a relationship) was associated with alcohol misuse.

No statistically significant interactions were found for employment, though for both samples, participants unable to work,

because they were long-term ill or disabled, were more likely to misuse alcohol (compared to those in employment).

### Sensitivity analysis

Results of our sensitivity analysis showed that male veterans were significantly more likely than non-veterans of the same age to exhibit probable CMD and alcohol misuse regardless of i) whether or not they had deployed and ii) whether or not they were deployed in a combat role (see online Supplementary Table S1).

When analyses were restricted only to veterans who had deployed, we found male veterans were significantly more likely than non-veterans of the same age to meet criteria for PTSD caseness. However, when comparing male veterans who had not been deployed with non-veterans of the same age, we found no significant difference between the two groups in the proportion of those who met criteria for probable PTSD. Male veterans were also significantly more likely than non-veterans to meet criteria for PTSD caseness when restricted to veterans deployed in a combat role. Prior to adjusting for education and marital status, we found no difference between veterans who were deployed in a non-combat role and non-veterans of the same age. After making this adjustment however, male veterans were significantly more likely to meet criteria for PTSD than non-veterans of the same age.

We found no statistically significant differences between female veterans and non-veterans when accounting for deployment.

### Discussion

This study has compared alcohol misuse, PTSD and CMD in a large representative sample of recent military service leavers to

**Table 4.** Regression analysis to show the association of socio-demographic characteristics on mental health for veterans and non-veterans separately

Socio-demographic Characteristics	Probable CMD						Probable PTSD						Probable Alcohol Misuse					
	KCMHR (n = 618)			UKHLS (n = 4063)			KCMHR (n = 187)			APMS (n = 293)			KCMHR (n = 264)			APMS (n = 175)		
	N	%	AOR (CI)	N	%	AOR (CI)	N	%	AOR (CI)	N	%	AOR (CI)	N	%	AOR	N	%	AOR (CI)
<b>Gender</b>																		
Male	557	23.3%	1	1466	15.5%	1	169	7.9%	1	101	5.0%	1	253	11.4%	1	106	6.1%	1
Female	61	23.1%	0.95 (0.67,1.36)	2597	21.2%	1.46 (1.35,1.58)	18	7.6%	0.86 (0.48,1.57)	192	6.8%	1.38 (1.03,1.85)	11	3.3%	0.25 (0.13,0.49)	69	2.2%	0.34 (0.24,0.49)
<b>Age</b>																		
22-34	122	27.3%	1	1048	17.1%	1	46	10.0%	1	84	5.0%	1	54	12.8%	1	46	6.4%	1
35-44	179	22.9%	0.79 (0.59,1.06)	854	15.5%	0.89 (0.77,1.03)	73	9.8%	0.98 (0.64,1.51)	61	5.3%	1.09 (0.62,1.90)	80	10.6%	0.82 (0.55,1.24)	46	6.7%	1.04 (0.62,1.75)
45-54	233	21.4%	0.73 (0.55,0.96)	994	16.8%	0.98 (0.85,1.13)	52	5.0%	0.47 (0.30,0.74)	71	6.2%	1.27 (0.74,2.18)	100	9.9%	0.72 (0.49,1.07)	39	5.3%	0.82 (0.47,1.41)
55-64	74	15.9%	0.50 (0.36,0.71)	772	16.0%	0.93 (0.80,1.08)	15	3.5%	0.32 (0.17,0.61)	57	6.5%	1.35 (0.79,2.30)	26	5.8%	0.40 (0.24,0.68)	32	3.8%	0.57 (0.32,1.03)
65-74	10	11.9%	0.36 (0.18,0.73)	395	9.5%	0.51 (0.42,0.62)	1	1.1%	0.10 (0.01,0.73)	20	1.2%	0.23 (0.10,0.55)	4	4.8%	0.33 (0.11,0.94)	12	1.5%	0.21 (0.09,0.48)
<b>Marital status</b>																		
Married/long-term relationship	489	20.9%	1	2513	14.2%	1	133	6.4%	1	94	2.8%	1	219	10.2%	1	72	3.9%	1
Single	55	34.3%	1.84 (1.22,2.77)	898	19.6%	1.45 (1.26,1.68)	21	13.5%	1.99 (1.11,3.55)	125	9.1%	3.78 (2.24,6.36)	18	11.8%	1.08 (0.58,2.00)	73	9.8%	2.71 (1.59,4.62)
Separated, divorced or widowed	69	39.2%	2.60 (1.76,3.83)	642	22.5%	1.83 (1.53,2.19)	30	18.7%	3.72 (2.16,6.41)	74	8.1%	2.84 (1.64,4.92)	26	17.0%	2.05 (1.18,3.58)	30	4.9%	1.57 (0.78,3.15)
<b>Highest qualification</b>																		
No qualifications	35	29.9%	1	437	20.7%	1	21	19.1%	1	101	12.5%	1	16	16.2%	1	35	7.2%	1
GCSE	138	23.9%	0.7 (0.41,1.21)	1037	16.7%	0.67 (0.54,0.83)	48	8.3%	0.36 (0.18,0.72)	89	5.0%	0.33 (0.19,0.57)	65	12.3%	0.72 (0.35,1.49)	57	6.9%	0.85 (0.45,1.61)
A level	201	23.5%	0.67 (0.40,1.15)	797	15.8%	0.63 (0.50,0.78)	62	8.2%	0.34 (0.17,0.67)	46	4.9%	0.32 (0.17,0.60)	91	11.1%	0.63 (0.31,1.27)	23	5.7%	0.65 (0.32,1.34)
Degree or higher	244	21.8%	0.67 (0.40,1.13)	1792	15.3%	0.60 (0.49,0.74)	56	5.6%	0.25 (0.13,0.51)	55	2.9%	0.18 (0.10,0.32)	92	8.2%	0.51 (0.26,1.01)	58	4.4%	0.51 (0.27,0.97)
<b>Employment status</b>																		
In employment (ft/pt)	507	21.5%	1	2329	13.3%	1	143	6.7%	1	100	3.2%	1	228	10.6%	1	103	5.1%	1
Unemployed (looking for work)	38	50.8%	3.88 (2.26,6.65)	365	32.3%	3.07 (2.51,3.75)	9	15.0%	2.46 (1.07,5.68)	22	11.6%	3.93 (1.65,9.36)	14	15.9%	1.58 (0.74,3.39)	15	13.4%	2.83 (1.29,6.21)
Long-term ill or disabled	52	79.5%	15.34 (7.69,30.58)	442	55.9%	8.56 (6.77,10.83)	31	52.8%	18.29 (9.83,34.05)	97	27.6%	11.38 (6.89,18.79)	13	19.6%	2.49 (1.22,5.07)	27	14.5%	3.52 (1.90,6.50)
Retired	14	8.5%	0.50 (0.27,0.93)	481	10.8%	1.18 (0.93,1.49)	2	1.0%	0.37 (0.10,1.42)	27	1.6%	0.87 (0.36,2.09)	8	5.1%	0.84 (0.36,1.96)	22	3.2%	3.12 (1.08,9.06)
Economically inactive	7	26.8%	1.28 (0.42,3.90)	446	24.0%	1.86 (1.45,2.38)	2	15.8%	2.63 (0.41,16.79)	47	8.4%	2.61 (1.34,5.10)	1	2.1%	0.21 (0.02,1.71)	8	2.8%	0.59 (0.17,2.12)
<b>NSSEC grade (employed only)</b>																		
1: Managerial and Professional	163	21.6%	1	1047	13.5%	1	43	5.5%	1	41	3.6%	1	75	9.4%	1	40	3.7%	1
2: Intermediate	107	23.3%	1.05 (0.74,1.49)	331	15.4%	1.13 (0.90,1.42)	32	6.7%	1.12 (0.61,2.07)	20	2.2%	0.56 (0.20,1.55)	37	9.4%	0.91 (0.54,1.54)	9	4.2%	1.16 (0.41,3.31)

3: Small and own employers	20	18.3%	0.76 (0.39,1.46)	197	12.4%	0.95 (0.75,1.19)	10	6.8%	1.09 (0.46,2.59)	14	2.8%	0.69 (0.26,1.83)	13	12.2%	1.17 (0.52,2.66)	18	7.6%	2.32 (1.03,5.23)
4: Lower supervisory & technical	32	20.6%	0.87 (0.50,1.49)	125	10.0%	0.72 (0.54,0.95)	13	9.8%	1.56 (0.66,3.66)	5	1.7%	0.46 (0.11,2.04)	21	16.8%	1.61 (0.84,3.07)	10	8.4%	2.28 (0.92,5.67)
5: Semi-routine/routine	110	21.6%	0.93 (0.66,1.32)	554	13.3%	0.97 (0.81,1.16)	51	10.9%	1.68 (0.95,2.98)	53	6.1%	1.6 (0.79,3.23)	47	11.8%	1.13 (0.70,1.83)	26	6.2%	1.65 (0.84,3.27)

AOR = adjusted for age and gender.

the general population. We found that, overall, UK veterans who served at the time of recent military operations were more likely to exhibit probable CMD, PTSD and alcohol misuse than non-veterans. These findings persist after adjusting for socio-demographic characteristics. When stratifying by working status, working veterans compared to working non-veterans were at a significant disadvantage for all three outcomes whereas non-working veterans compared to non-working non-veterans were more likely to report probable CMD and PTSD. Stratifying by gender showed that the negative impact of being a veteran on mental health and alcohol misuse was restricted to male veterans. Being ill or disabled was associated with CMD and PTSD for both veterans and non-veterans.

The only previous UK-wide mental health comparison study between veterans and non-veterans – Woodhead *et al.* (2011a, b) – found no significant difference between the two groups. Woodhead *et al.* (2011a, b) utilised a previous cohort of veterans, few of whom had served during recent military operations in Iraq and Afghanistan. This might suggest that the mental health disparity between veterans and non-veterans in the current study could have resulted from deployment to these recent operations. The findings of our sensitivity analysis partially support this hypothesis. They show that while veterans who had been deployed to Iraq or Afghanistan were more likely to meet criteria for probable PTSD than non-veterans of the same age, veterans who had not been deployed were not more likely to meet criteria. Male veterans were more likely to meet criteria for probable CMD and misuse of alcohol than non-veterans of the same age regardless of whether they were deployed to Iraq or Afghanistan, or whether they were deployed in a combat role. Deployment to Iraq or Afghanistan therefore may not be the only factor contributing to the increased levels of CMDs and alcohol misuse in male veterans. It is possible that other vulnerability factors, such as childhood adversity (Iversen *et al.*, 2007), family separations, work-related injuries, struggling to adapt to civilian life may also explain the difference for CMD and alcohol use (the latter may also reflect a continuation of heavy drinking from military life). Alternatively, our contradictory findings to Woodhead *et al.* (2011b) could be due to different age composition of the sample used in this study or recent changes made to the process of leaving the military.

There are notable gender differences in our findings. Among non-veterans, CMD and PTSD prevalence were greater in women compared to men [in line with recent estimates (McManus *et al.*, 2016)]. For veterans, the prevalence of CMD and PTSD was significantly higher than non-veterans for men only. This is partially supported by some PTSD studies which have found no gender differences in military and police samples (Pole *et al.*, 2001; Sutker, Davis, Uddo, & Ditta, 1995). However, the finding that female veterans are no more likely to have CMD than male veterans contradicts USA and previous UK studies (Haskell *et al.*, 2010; Stansfeld, Rasul, Head, & Singleton, 2011). Overall, for CMD and PTSD, caseness was generally much lower in working people (veterans and non-veterans) compared to non-working people, and specifically in working veterans compared to non-working non-veterans.

Both male and female veterans are more likely to report symptoms of drinking at a hazardous level than their non-veteran counterparts. Furthermore, male veterans are more likely to report symptoms of dependence and harmful drinking than men of the same age. These findings, combined with previous studies, indicate that a culture of excessive and persistent drinking



exists in the military compared to the general population (Fear *et al.*, 2007; Goodwin *et al.*, 2017). This suggests that drinking habits acquired before or during military service endure after transitioning into non-veteran life. A recent UK study has also found hazardous drinking to be prevalent amongst female spouses of military personnel (Gribble, Goodwin, & Fear, 2019), suggesting that there are implications of the drinking culture in females both within and connected to the AF. This is important as evidence suggests that women may be particularly susceptible to alcohol-related physical and mental harm (Erol & Karpyak, 2015). Working and non-working veterans report similarly high levels of alcohol misuse. Unlike in the non-veteran sample, employment is not associated with reduced alcohol misuse in veterans. Further work is needed to understand this finding, such as exploration of the type of work undertaken by veterans compared to non-veterans.

Unemployed and ill or disabled veterans were more likely to exhibit probable CMD and PTSD than veterans in employment. This is supported by previous research which has found poor mental health to be high in unemployed veterans compared to those in employment (Hamilton, Williams, & Washington, 2015; Harden & Murphy, 2018). A high prevalence of CMD has also been identified among veterans with a physical impairment (Stevellink *et al.*, 2015). Further longitudinal analysis is needed, not only to unpick the causal relationship between employment and mental health status for veterans but to determine if physical health affects veteran's ability to work and whether this in turn negatively impacts their mental health.

### Implications

Findings from this study demonstrate that veterans (especially male veterans) are at a disadvantage to non-veterans in respect of their mental health. In particular, it appears that service leavers may benefit from interventions targeting alcohol misuse, with recent evidence suggesting that electronic interventions could be suitable (Leightley, Puddephatt, Goodwin, Rona, & Fear, 2018). This is especially important for those who had harmful and persistent drinking habits during their military service. Veterans who are unable to work for health reasons may also benefit from additional social support. Being removed from the support and structure of military life and unable to work, they are at risk of becoming isolated which in turn can negatively impact their health (Elovainio *et al.*, 2017; Hatch *et al.*, 2013).

### Strengths and limitations

This study utilises data representative of veterans who served in the UK military during recent military operations in Afghanistan and Iraq, allowing us to gain an insight into the mental health of recent service leavers compared to non-veterans.

This study does have some limitations. Results are derived from self-reported data and geographic distribution of participants could not be accounted for. Many of the veterans who participated in phase 3 of the KCMHR cohort left service around the time of their participation in phase 1 (2004–2006), the results of our study are therefore not necessarily representative of veterans who left service more than 12 years ago. These veterans also are more likely to be aware of the most common symptoms of PTSD than civilians due to ongoing efforts in the military to

increase awareness and reduce stigma, which in some cases could have resulted in veterans being more likely to have reported symptoms. We acknowledge this study is restricted to using DSM-IV criteria to assess probable PTSD domain caseness, in order to use comparable measures across populations. The general population surveys utilised in this survey used different methods of data collection than the KCMHR study. APMS was conducted in England only while UKHLS and KCMHR were conducted UK-wide. Due to the lack of available data in the non-veteran population, we were also unable to examine the potential moderating effect of childhood adversity on the association between military service and mental health. Furthermore, due to the small number of older veterans, entropy balancing may not have been fully able (in some cases) to resolve the differences between the samples. Finally, our veteran sample is only 10% female ( $n = 305$ ), and whilst this is representative of the UK veteran population, some of our analysis restricted to females may be underpowered.

### Conclusion

It is only by understanding the relative mental health burden experienced by veterans that the Government can ensure that they are at 'no disadvantage' which is a key principle of the AF covenant (Taylor, 2011). Our findings demonstrate that, whilst the same sociodemographic groups within the veteran and non-veteran populations seemed to have an increased risk of mental health problems (e.g. those who were unemployed), male veterans, in particular, appear to be at a distinct disadvantage compared to those who have never served.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291720001944>.

**Author contributions.** DM and LG conceived the study concept and design with input from NF and NG. RR, DM and LG contributed to the design of the analysis. MJ curated the military dataset utilised in this study. RR conducted the statistical analysis and wrote the article supervised by DM and LG. All authors provided feedback on the manuscript.

**Acknowledgements.** We would like to thank the Forces in Mind Trust for funding this study (FiMT16/0304K).

The KCMHR study was funded by the UK Ministry of Defence.

APMS 2014 was commissioned by NHS Digital, funded by the Department of Health, and carried out by NatCen Social Research and the University of Leicester.

UKHLS is primarily funded by the Economic Social Research Council but is also supported both financially and in kind by a wide range of other co-funding organisations detailed here: <https://www.understandingsociety.ac.uk/about/funders>

The funders of the study had no role in study design, data analysis, or data interpretation. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication. Forces in Mind Trust (FiMT16/0304 K).

**Conflict of interest.** NTF is a trustee of a charity providing support to veterans and their families and is a member of IGARD. DM is a Lead Consultant Psychiatrist for an NHS veteran mental health service for London and the South East of England. NG is the Royal College of Psychiatrists lead for military health.

**Ethical standards.** This study received ethical approval from the Proportionate Review Sub-committee of the West Midlands-- South Birmingham Research Ethics Committee (REC reference:17/WM/0147, IRAS project ID: 224664).

Ethical approval for the KCMHR study was granted by the UK Ministry of Defence Research Ethics Committee (reference: 448/MODREC/13) and the

King's College London Psychiatry Nursing and Midwifery Research Ethics Subcommittee (reference: PNM/12/13–169).

Ethical approval for the ongoing UKHLS survey was obtained from University of Essex Ethics Committee and is detailed in the UKHLS user guide, available at: <https://www.understandingsociety.ac.uk/sites/default/files/downloads/documentation/mainstage/user-guides/mainstage-user-guide.pdf>

Ethical approval for APMS 2014 was obtained from the West London National Research Ethics Committee (reference: 14/LO/0411)

## References

- Babor, T. F., Higgins-Biddle, J. C., Saunders, J. B., & Monteiro, M. G., & World Health Organization (2001). AUDIT: The alcohol use disorders identification test: Guidelines for use in primary health care.
- Defence Select Committee (2018). Mental health and the Armed Forces, Part One: The Scale of mental health issues: Government Response to the Committee's Eleventh Report. (HC 813).
- Elovainio, M., Hakulinen, C., Pulkki-Råback, L., Virtanen, M., Josefsson, K., Jokela, M., ... Kivimäki, M. (2017). Contribution of risk factors to excess mortality in isolated and lonely individuals: An analysis of data from the UK Biobank cohort study. *The Lancet Public Health*, 2(6), e260–e266. doi:10.1016/S2468-2667(17)30075-0.
- Erol, A., & Karpyak, V. M. (2015). Sex and gender-related differences in alcohol use and its consequences: Contemporary knowledge and future research considerations. *Drug and Alcohol Dependence*, 156, 1–13. doi:10.1016/j.drugalcdep.2015.08.023.
- Fear, N. T., Iversen, A., Meltzer, H., Workman, L., Hull, L., Greenberg, N., ... Wessely, S. (2007). Patterns of drinking in the UK Armed Forces. *Addiction*, 102(11), 1749–1759. doi:10.1111/j.1360-0443.2007.01978.x.
- Fear, N. T., Jones, M., Murphy, D., Hull, L., Iversen, A. C., Coker, B., ... Wessely, S. (2010). What are the consequences of deployment to Iraq and Afghanistan on the mental health of the UK armed forces? A cohort study. *The Lancet*, 375(9728), 1783–1797. doi:10.1016/S0140-6736(10)60672-1.
- Goldberg, D. P., Gater, R., Sartorius, N., Ustun, T. B., Piccinelli, M., Gureje, O., & Rutter, C. (1997). The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychological Medicine*, 27(1), 191–197.
- Goodwin, L., Norton, S., Fear, N. T., Jones, M., Hull, L., Wessely, S., & Rona, R. J. (2017). Trajectories of alcohol use in the UK military and associations with mental health. *Addictive Behaviors*, 75, 130–137. doi:10.1016/j.addbeh.2017.07.010.
- Goodwin, L., Wessely, S., Hotopf, M., Jones, M., Greenberg, N., Rona, R. J., ... Fear, N. T. (2015). Are common mental disorders more prevalent in the UK serving military compared to the general working population? *Psychological Medicine*, 45(9), 1881–1891. doi:10.1017/S0033291714002980.
- Gribble, R., Goodwin, L., & Fear, N. T. (2019). Mental health outcomes and alcohol consumption among UK military spouses/partners: A comparison with women in the general population. *European Journal of Psychotraumatology*, 10(1). <https://doi.org/10.1080/2008198.2019.1654781>.
- Hainmueller, J. (2012). Entropy balancing for causal effects: A multivariate reweighting method to produce balanced samples in observational studies. *Political Analysis*, 20(1), 25–46.
- Hamilton, A. B., Williams, L., & Washington, D. L. (2015). Military and mental health correlates of unemployment in a national sample of women veterans. *Medical Care*, 53, S32. doi:10.1097/MLR.0000000000000297.
- Harden, L., & Murphy, D. (2018). Risk factors of suicidal ideation in a population of UK military veterans seeking support for mental health difficulties. *Journal of the Royal Army Medical Corps*, 164(5), 352–356. doi:10.1136/jramc-2018-000921.
- Haskell, S. G., Gordon, K. S., Mattocks, K., Duggal, M., Erdos, J., Justice, A., & Brandt, C. A. (2010). Gender differences in rates of depression, PTSD, pain, obesity, and military sexual trauma among Connecticut War Veterans of Iraq and Afghanistan. *Journal of Women's Health* (2002), 19(2), 267–271. doi:10.1089/jwh.2008.1262.
- Hatch, S. L., Harvey, S. B., Dandeker, C., Burdett, H., Greenberg, N., Fear, N. T., & Wessely, S. (2013). Life in and after the Armed Forces: Social networks and mental health in the UK military. *Sociology of Health & Illness*, 35(7), 1045–1064. doi:10.1111/1467-9566.12022.
- Hoerster, K. D., Lehavot, K., Simpson, T., McFall, M., Reiber, G., & Nelson, K. M. (2012). Health and health behavior differences: U.S. Military, Veteran, and Civilian Men. *American Journal of Preventive Medicine*, 43(5), 483–489. doi:10.1016/j.amepre.2012.07.029.
- Hooff, M. V., McFarlane, A. C., Davies, C. E., Searle, A. K., Fairweather-Schmidt, A. K., Verhagen, A., ... Hodson, S. E. (2014). The Australian defence force mental health prevalence and wellbeing study: Design and methods. *European Journal of Psychotraumatology*, 5(1), 23950. doi:10.3402/ejpt.v5.23950.
- Hotopf, M., Hull, L., Fear, N. T., Browne, T., Horn, O., Iversen, A., ... Wessely, S. (2006). The health of UK military personnel who deployed to the 2003 Iraq war: A cohort study. *Lancet (London, England)*, 367(9524), 1731–1741. doi:10.1016/S0140-6736(06)68662-5.
- Iversen, A. C., Fear, N. T., Simonoff, E., Hull, L., Horn, O., Greenberg, N., ... Wessely, S. (2007). Influence of childhood adversity on health among male UK military personnel. *The British Journal of Psychiatry*, 191(6), 506–511. doi:10.1192/bjp.bp.107.039818.
- Iversen, A. C., & Greenberg, N. (2009). Mental health of regular and reserve military veterans. *Advances in Psychiatric Treatment*, 15(2), 100–106. doi:10.1192/apt.bp.107.004713.
- Lee, J. E., Sudom, K. A., & Zamorski, M. A. (2013). Longitudinal analysis of psychological resilience and mental health in Canadian military personnel returning from overseas deployment. *Journal of Occupational Health Psychology*, 18(3), 327.
- Leightley, D., Puddephatt, J.-A., Goodwin, L., Rona, R., & Fear, N. T. (2018). InDEx: Open source iOS and android software for self-reporting and monitoring of alcohol consumption. *Journal of Open Research Software*, 6, 13–24. doi:10.5334/jors.207
- Lynn, P. (2006). *Quality profile: BHPS version 2.0: Waves 1 to 13 1991–2003*. Essex, UK: University of Essex. Institute for Social Research.
- McManus, S., Bebbington, P., Jenkins, R., & Brugha, T. (2016). *Mental health and wellbeing in England: Adult psychiatric morbidity survey 2014*. Leed, UK: NHS Digital.
- Murphy, D., Palmer, E., Westwood, G., Busuttill, W., & Greenberg, N. (2016). Do alcohol misuse, service utilisation, and demographic characteristics differ between UK veterans and members of the general public attending an NHS general hospital? *Journal of Clinical Medicine*, 5(11), 95–107. doi:10.3390/jcm5110095
- Office for National Statistics (n.d.). NS-SEC. The National Statistics Socio-Economic Classification (NS-SEC) - Office for National Statistics. Retrieved 6 March 2018, from <https://www.ons.gov.uk/methodology/classificationsandstandards/otherclassifications/thenationalstatisticsocioeconomicclassificationnssecbasedonsoc2010>.
- Oster, C., Morello, A., Venning, A., Redpath, P., & Lawn, S. (2017). The health and wellbeing needs of veterans: A rapid review. *BMC Psychiatry*, 17(1), 411–428. doi:10.1186/s12888-017-1547-0
- Ostler, K., Thompson, C., Kinmonth, A.-L. K., Peveler, R. C., Stevens, L., & Stevens, A. (2001). Influence of socio-economic deprivation on the prevalence and outcome of depression in primary care: The Hampshire Depression Project. *The British Journal of Psychiatry*, 178(1), 12–17. doi:10.1192/bjp.178.1.12.
- Pole, N., Best, S. R., Weiss, D. S., Metzler, T., Liberman, A. M., Fagan, J., & Marmar, C. R. (2001). Effects of gender and ethnicity on duty-related post-traumatic stress symptoms among urban police officers. *The Journal of Nervous and Mental Disease*, 189(7), 442–448.
- Royal British Legion (2014). A UK household survey of the ex-service community.
- Stansfeld, S. A., Rasul, F. R., Head, J., & Singleton, N. (2011). Occupation and mental health in a national UK survey. *Social Psychiatry and Psychiatric Epidemiology*, 46(2), 101–110. doi:10.1007/s00127-009-0173-7.
- Stevellink, S., Jones, M., Hull, L., Pernet, D., MacCrimmon, S., Goodwin, L., ... Wessely, S. (2018). Mental health outcomes at the end of the British involvement in the Iraq and Afghanistan conflicts: A cohort study. *The British Journal of Psychiatry: The Journal of Mental Science*, 213(6), 690–697. doi:10.1192/bjp.2018.175
- Stevellink, S., Malcolm, E. M., Mason, C., Jenkins, S., Sundin, J., & Fear, N. T. (2015). The prevalence of mental health disorders in (ex-)military personnel with a physical impairment: A systematic review. *Occupational and Environmental Medicine*, 72(4), 243–251. doi:10.1136/oemed-2014-102207.

- Sundin, J., Herrell, R. K., Hoge, C. W., Fear, N. T., Adler, A. B., Greenberg, N., ... Bliese, P. D. (2014). Mental health outcomes in US and UK military personnel returning from Iraq. *The British Journal of Psychiatry*, 204(3), 200–207. doi:10.1192/bjp.bp.113.129569.
- Sutker, P. B., Davis, J. M., Uddo, M., & Ditta, S. R. (1995). Assessment of psychological distress in Persian gulf troops: Ethnicity and gender comparisons. *Journal of Personality Assessment*, 64(3), 415–427. doi:10.1207/s15327752jpa6403\_2.
- Taylor, C. (2011). *Armed forces covenant. Standard note SN/IA/5979*. London: Government Publication.
- Taylor, S., Koch, W. J., Kuch, K., Crockett, D. J., & Passey, G. (1998). The structure of posttraumatic stress symptoms. *Journal of Abnormal Psychology*, 107(1), 154–160. Scopus. doi:0.1037/0021-843X.107.1.154.
- Thandi, G., Sundin, J., Ng-Knight, T., Jones, M., Hull, L., Jones, N., ... Fear, N. T. (2015). Alcohol misuse in the United Kingdom Armed Forces: A longitudinal study. *Drug and Alcohol Dependence*, 156, 78–83. doi:10.1016/j.drugalcdep.2015.08.033.
- Vasterling, J. J., Proctor, S. P., Friedman, M. J., Hoge, C. W., Heeren, T., King, L. A., & King, D. W. (2010). PTSD Symptom increases in Iraq-deployed soldiers: Comparison with nondeployed soldiers and associations with baseline symptoms, deployment experiences, and postdeployment stress. *Journal of Traumatic Stress*, 23(1), 41–51. doi:10.1002/jts.20487.
- Weathers, F. W., Litz, B. T., Herman, D., Huska, J., & Keane, T. (1994). *The PTSD checklist-civilian version (PCL-C)*. Boston, MA: National Center for PTSD.
- Wilkins, K. C., Lang, A. J., & Norman, S. B. (2011). Synthesis of the psychometric properties of the PTSD checklist (PCL) military, civilian, and specific versions. *Depression and Anxiety*, 28(7), 596–606. doi:10.1002/da.20837.
- Woodhead, C., Rona, R. J., Iversen, A., MacManus, D., Hotopf, M., Dean, K., ... Fear, N. T. (2011b). Mental health and health service use among post-national service veterans: Results from the 2007 Adult Psychiatric Morbidity Survey of England. *Psychological Medicine*, 41(2), 363–372. doi:10.1017/S0033291710000759.
- Woodhead, C., Rona, R. J., Iversen, A. C., MacManus, D., Hotopf, M., Dean, K., ... Fear, N. T. (2011a). Health of national service veterans: An analysis of a community-based sample using data from the 2007 Adult Psychiatric Morbidity Survey of England. *Social Psychiatry and Psychiatric Epidemiology*, 46(7), 559–566. doi:10.1007/s00127-010-0232-0.